International Journal of Advance Research in Science, Management, Engineering and Technology
Volume X, Issue X | Month Year | www.ijarsmet.com
ISSN: 3139-1141

Environmental effects of human activities: a review of agriculture, housing, industry, mining and transportation

Sectoral pressures, cumulative impacts and integrated mitigation

Sushma M.V., Likhita C.D., Lokesh M.M., Gagan D.S., Anjali Sharma

Sushma M.V. (Assistant Professor), Department of Computer Science and Engineering (Artificial Intelligence and Machine Learning), Malnad College of Engineering, Hassan, Karnataka, India

Abstract—Human activities provide food, shelter, manufactured goods, minerals and mobility, but also place substantial pressure on climate, ecosystems, freshwater and human health. This review examines the environmental effects of agriculture, housing and construction, industry, mining and transportation through their principal pathways: greenhouse-gas emissions, land-cover change, soil degradation, water use and contamination, air pollution, waste and biodiversity loss. A narrative synthesis of major international assessments and institutional literature was used, with emphasis on the Intergovernmental Panel on Climate Change, the Food and Agriculture Organization of the United Nations, the United Nations Environment Programme and the World Health Organization. The evidence indicates that impacts are interconnected: housing demand stimulates material extraction and industrial production; transport infrastructure changes land use and access; agricultural supply chains depend on energy and logistics; and mining can create persistent landscape and water risks while supplying materials used by other sectors. The paper compares sector-specific pressures and discusses responses including soil and nutrient management, compact urban planning, low-carbon construction, cleaner industrial production, responsible mineral governance, public transport, electrification and circular resource use. It argues that durable environmental improvement requires life-cycle assessment, cumulative-impact management, transparent monitoring and equitable participation. This is a literature-based review and does not report new field experiments or a quantitative meta-analysis.

Keywords—human activities; environmental degradation; agriculture; housing; industry; mining; transportation; climate change; biodiversity; pollution; sustainability

I. INTRODUCTION

Human societies alter natural systems to secure food, shelter, energy, materials and mobility. These activities have supported health, trade and economic development, but their scale and design can exceed the regenerative capacity of ecosystems. Environmental consequences include climate change, habitat conversion, soil erosion, freshwater stress, chemical contamination, air pollution, waste accumulation and unequal exposure to environmental hazards. The Intergovernmental Panel on Climate Change (IPCC) concludes that human activities, principally through greenhouse-gas emissions, have unequivocally caused global warming [1]. Its assessments also show that mitigation depends on coordinated action across energy, industry, transport, buildings and land-use systems [2].

Agriculture, housing and construction, industry, mining and transportation are linked rather than independent. Housing uses cement, steel, timber and minerals; industry processes raw materials and supplies equipment; transport moves goods and shapes urban form; agriculture relies on energy, fertilizer and logistics; and mining supplies materials for both conventional infrastructure and low-emission technologies. A narrow sector-by-sector analysis may therefore miss indirect impacts and trade-offs. This review examines each activity, compares their principal environmental pathways and identifies responses that protect ecosystems while supporting human well-being.

The contribution of this review is an integrated comparison of the five activities through shared environmental receptors and decision points. Rather than ranking sectors with one unsupported universal score, the paper distinguishes pressures, local exposure, life-cycle effects and management options. This is important because the same activity can have very different consequences depending on ecosystem sensitivity, technology, governance and the distribution of costs and benefits.

A. Aim and objectives

The aim is to critically synthesize evidence on the environmental effects of five major human activities and identify practical strategies for reducing their impacts.

- Identify the main effects of agriculture, housing, industry, mining and transportation on air, water, land, climate and biodiversity.
- Summarize findings from authoritative international assessments and institutional literature.
- Compare shared impacts, sector interactions and potential trade-offs.
- Recommend prevention, mitigation, restoration and governance measures that can be applied across sectors.

ISSN: 3139-1141

II. REVIEW METHODOLOGY

A narrative literature-review approach was used to synthesize major scientific assessments and institutional publications. Priority was given to the IPCC Sixth Assessment Report, the Food and Agriculture Organization of the United Nations (FAO) assessment of land and water resources, United Nations Environment Programme (UNEP) publications on buildings, pollution and mining-related biodiversity risks, and World Health Organization (WHO) material on transport-related air pollution [1]–[7]. These sources were selected for their broad geographic scope, transparent institutional authorship and relevance to the five activity domains.

The review was organized around five sectors and seven environmental pathways: greenhouse-gas emissions, land transformation, soil condition, water quantity and quality, air pollution, waste and biodiversity. Cross-sector synthesis was used to identify interactions, justice concerns and opportunities for coordinated policy. This paper is not a preregistered systematic review, a meta-analysis or an original field study. It does not claim to cover every publication worldwide, and it reports no new experiments, interviews or measurements. Statistics from different reports may use different years, boundaries and accounting conventions; they are therefore interpreted in context rather than combined as if directly comparable.

Evidence was prioritized when it came from global scientific assessments, official technical reports or authoritative health and environmental agencies. Findings are presented qualitatively because the selected sources use different reference years, spatial scales and system boundaries. No numerical meta-analysis was performed, and no causal estimate is attributed to this review. The reference list is intentionally traceable so readers can consult the original assessments and judge their applicability to local conditions.

III. LITERATURE REVIEW

A. *Climate change and land use*

The IPCC assesses greenhouse-gas emissions across energy, industrial processes, transport, buildings and agriculture, forestry and other land use. The literature emphasizes that emissions depend on technology, infrastructure, demand, land management and consumption patterns, not simply on the number of individual devices or facilities [1], [2]. Land-based mitigation can include protecting forests, peatlands and wetlands, improving soil management and reducing food loss, but poorly designed interventions may compete with food production or damage biodiversity.

B. *Land, soil and freshwater*

FAO's global assessment highlights the importance of sustainable soil and water management in food systems. Its 2025 report states that more than 60% of human-induced land degradation occurs on agricultural land and that agriculture accounts for more than 70% of global freshwater withdrawals [3]. These findings support integrated soil conservation, efficient irrigation, watershed management and locally appropriate agricultural practices.

C. *Buildings, industry and material demand*

UNEP and the Global Alliance for Buildings and Construction report that buildings and construction have a substantial energy, emissions and material footprint. The response agenda includes improved building codes, efficient design, low-carbon materials, renewable energy and circular construction [4]. For heavy industry, the IPCC identifies energy and material efficiency, electrification, low-emissions fuels, alternative feedstocks and selected carbon-capture applications as possible routes to lower emissions, while noting that technical feasibility differs by process [2].

A newer global assessment for 2025–2026 reports that buildings and construction account for approximately 37% of global emissions and nearly half of global material extraction under its stated accounting boundaries [8]. These figures should not be directly substituted for older estimates without checking definitions: reports may include different combinations of building operations, construction materials and infrastructure. The important conclusion is consistent across assessments: both operational energy and embodied materials need attention.

D. *Mining, pollution and biodiversity*

UNEP's work on artisanal and small-scale gold mining documents risks associated with land disturbance, chemical use and contamination when extraction is poorly managed [5]. More broadly, mining impacts depend on geology, processing methods, hydrology, tailings design, site location and the quality of closure planning. UNEP also identifies chemical pollution as a significant pressure on ecosystems and biodiversity [6].

E. *Transportation and health*

The IPCC frames transport mitigation as a system question as well as a vehicle-technology question. Urban form, trip distances, public transport availability, freight organization and demand for mobility influence the scale of energy use before a vehicle technology is selected [2]. This supports the avoid–shift–improve framework used later in the paper.

WHO describes transport-related air pollution as including both exhaust emissions and non-exhaust particles from brakes, tyres and resuspended road dust. Exposure to these pollutants is associated with adverse health outcomes, including cardiovascular disease and asthma [7]. Transport policy therefore needs to consider climate emissions alongside local air quality, noise, road safety, access and the distribution of exposure.

IV. ENVIRONMENTAL EFFECTS OF AGRICULTURE

A. *Land conversion and habitat loss*

Agricultural expansion can also affect ecosystems indirectly through feed production, commodity demand, water infrastructure and new access roads. Land-use decisions should therefore assess both the farm plot and the wider supply chain. Protecting intact ecosystems is often more effective than attempting to recreate their full ecological complexity after conversion.

Agriculture requires land for crops, livestock and supporting infrastructure. Expansion into forests, wetlands, grasslands and other natural habitats removes vegetation, reduces carbon storage and fragments ecosystems. The magnitude of the impact depends on the original habitat, the area converted, production practices and the supply-chain drivers of expansion. Agricultural intensification on existing

 ISSN: 3139-1141

farmland may reduce pressure for conversion in some settings, but only if productivity gains are not accompanied by greater total demand or damaging input use.

B. *Soil degradation and erosion*

Soil health is multidimensional: organic matter, structure, infiltration, nutrient balance, biological activity and erosion resistance all influence long-term productivity. A practice that improves one measure may not improve every measure in every climate. Soil-cover and tillage decisions should be evaluated against local rainfall, slope, crop residues, fuel use and soil type rather than promoted as a universal prescription.

Intensive tillage, overgrazing, removal of crop residues, cultivation on steep slopes and poorly managed irrigation can accelerate erosion and reduce soil organic matter. Sediment may carry nutrients and pesticides into waterways, while compaction reduces infiltration and root growth. In irrigated drylands, inadequate drainage can contribute to salinity. Soil degradation may reduce yields and encourage additional inputs or land conversion, creating a reinforcing cycle of environmental pressure.

C. *Water use, nutrient runoff and chemicals*

Nutrient management is most effective when source, rate, timing and placement are considered together. Riparian buffers and constructed wetlands can intercept some runoff, but they are not substitutes for preventing excessive application at source. Basin-level governance is needed where irrigation users share rivers or aquifers, because farm-level efficiency gains do not always reduce total withdrawals if saved water is used to expand irrigated area.

Irrigation is essential in many regions, but withdrawals that exceed recharge or ecological-flow requirements can deplete rivers and groundwater. Nitrogen and phosphorus from fertilizers and manure can enter water bodies and promote eutrophication. Pesticides may harm aquatic organisms and non-target species, while untreated farm effluent can increase pathogen loads. Risks can be reduced through soil testing, nutrient budgeting, precise application, integrated pest management, riparian buffers and irrigation schedules matched to crop demand and local water availability.

D. *Climate and biodiversity impacts*

Climate resilience and biodiversity protection can reinforce each other when practices maintain soil moisture, landscape diversity and habitat connectivity. However, land-based climate interventions should be checked for trade-offs, including water demand, competition with food production and the use of non-native species. Emissions reductions should be measured against a clear baseline and should not be assumed from the adoption of a practice alone.

Agriculture emits methane from ruminant digestion and some rice systems, nitrous oxide from soils and nitrogen inputs, and carbon dioxide from fuel use and land-use change. Farm landscapes simplified into large monocultures may provide fewer resources for pollinators, birds, natural pest predators and soil organisms. Practices such as crop rotation, agroforestry, habitat strips, improved manure management and protection of native vegetation can support biodiversity and reduce some emissions, provided they are adapted to local conditions.

E. *Sustainable agricultural responses*

- Maintain soil cover and use erosion-control practices appropriate to slope, soil and climate.
- Apply fertilizers according to crop demand and soil tests, and avoid application when heavy rainfall is forecast.
- Use integrated pest management and targeted pesticide application to limit unnecessary chemical exposure.
- Improve irrigation scheduling and manage groundwater withdrawals within sustainable limits.
- Protect remaining high-value ecosystems and reduce food loss and waste throughout supply chains.




Fig. 1. Agricultural pressure pathways and practical intervention points.

- Promote locally suitable crop diversity, agroforestry and climate-resilient production.

V. ENVIRONMENTAL EFFECTS OF HOUSING AND CONSTRUCTION

A. *Land take and urban expansion*

Urban land-use decisions can lock in environmental outcomes for decades because streets, buildings and utility networks are long-lived assets. Infill development and transit-oriented growth may reduce land take and trip distances, but only if housing remains accessible and urban green space, drainage and heat mitigation are protected. Planning should also consider informal settlements and avoid pushing lower-income residents into distant locations with poor transport access.

Housing development converts land to buildings, roads, parking, utilities and other infrastructure. Unplanned expansion can fragment habitats, replace farmland and wetlands, increase runoff and place pressure on peri-urban water supplies. Low-density development can also lengthen travel distances and increase dependence on private vehicles. Construction on floodplains, steep slopes or coastlines may expose people and ecosystems to hazards and raise future adaptation costs.

B. *Embodied carbon and material extraction*

Embodied impacts include raw-material extraction, processing, manufacturing, transport, construction, maintenance and end-of-life treatment. Comparing materials requires a functional unit and equivalent performance, durability, fire safety and maintenance assumptions. Environmental product declarations and life-cycle assessment can support decisions, but results should be checked for consistent boundaries and local electricity mixes.

ISSN: 3139-1141

A building's footprint begins before occupation. Cement, steel, bricks, glass, aluminium, timber and insulation require extraction, manufacture and transport. Cement production releases carbon dioxide from fuel use and from limestone calcination, while steelmaking can be highly energy-intensive. Poorly managed sand, gravel and timber extraction can damage river systems and forests. Structural efficiency, reuse, recycled content and verified lower-carbon materials can reduce demand for virgin resources.

C. *Operational energy, water and waste*

Operational performance is shaped by climate, occupant behavior, building controls, maintenance and the carbon intensity of electricity. A building designed efficiently can still perform poorly if systems are badly commissioned or maintained. Water-sensitive design should account for local rainfall, water quality, public-health requirements and the reliability of municipal supply; rainwater reuse must be designed for its intended purpose.

Buildings use energy for cooling, heating, ventilation, lighting, appliances and water heating. Climate-responsive design, shading, insulation and efficient equipment can reduce demand. Water is required for drinking, sanitation and landscaping; leaks and inefficient fixtures waste treated supplies. Impervious surfaces can increase flood peaks and transport contaminants into waterways. Construction and demolition also generate concrete, timber, metals, glass, packaging and potentially hazardous materials. Waste separation, safe handling and material recovery can reduce disposal and virgin-material demand.

D. *Sustainable housing responses*

- Encourage compact, mixed-use development near reliable public transport while protecting floodplains and ecological corridors.
- Prioritize passive cooling, daylight, ventilation and climate-appropriate building envelopes.
- Use efficient appliances, low-emissions electricity and rooftop solar where feasible.
- Design buildings for durability, repair, adaptive reuse and efficient material use.
- Separate construction waste and reuse or recycle suitable concrete, metal, timber and glass.




Fig. 2. Whole-life stages of housing and the principal opportunities to reduce environmental burden.

- Install water-efficient fixtures and sustainable drainage measures where locally appropriate.

VI. ENVIRONMENTAL EFFECTS OF INDUSTRY

A. *Greenhouse gases and air pollution*

Industrial emissions can be concentrated around facilities and supply chains, making both regional climate accounting and local exposure assessment necessary. Continuous monitoring, stack testing, fugitive-emission surveys and worker exposure measurements answer different questions and should not be treated as interchangeable. Emission limits are most effective when accompanied by maintenance requirements, inspections, credible penalties and public reporting.

Industrial impacts arise from fuel combustion and chemical processes. Steel, cement, chemicals, glass, pulp and paper and other manufacturing sectors may require substantial heat and electricity. Some processes release carbon dioxide even when energy efficiency improves. Industrial facilities can also emit particulate matter, sulfur dioxide, nitrogen oxides, volatile organic compounds and toxic metals. These pollutants can affect workers and nearby communities, contribute to secondary air pollution and deposit onto land and water.

B. *Water use, wastewater and hazardous waste*

Wastewater controls should follow a pollutant-specific risk assessment rather than a single generic treatment assumption. Some contaminants require source substitution or specialized treatment, and conventional treatment may not remove all persistent chemicals or dissolved metals. Safe waste management also requires traceability from generation to transport, treatment and final disposal, with particular attention to accidental releases and legacy contamination.

Water is used for cooling, cleaning, processing and manufacturing. Depending on the sector, effluent can contain organic matter, salts, acids, dyes, oils, solvents, nutrients or metals. Inadequate treatment can damage aquatic ecosystems and compromise downstream uses. Industrial waste streams may include hazardous sludge, solvents, ash, chemical residues and electronic waste. Pollution prevention, effective treatment, secure storage and transparent monitoring are essential, because control systems only work when maintained and operated properly.

C. *Cleaner industrial production*

Industrial mitigation requires process-specific solutions. Measures include energy audits, efficient motors, heat recovery, electrification of suitable processes, low-emissions electricity, alternative feedstocks, material efficiency and product redesign. Industrial symbiosis can allow a safe by-product from one facility to replace a raw material elsewhere. Circular approaches can reduce demand for virgin resources, but recycling must also manage energy use and potential toxic exposures. Emission and effluent data should be measured consistently and reported with clear boundaries.

ISSN: 3139-1141




Fig. 3. A pollution-prevention hierarchy for industrial facilities.

VII. ENVIRONMENTAL EFFECTS OF MINING

A. *Landscape and ecosystem disturbance*

The mine footprint includes more than the excavation itself: waste-rock areas, tailings, haul roads, power lines, water diversions and worker settlements can expand the affected landscape. Environmental impact assessment should map sensitive habitats, seasonal species use, groundwater dependence and cumulative pressures from other projects in the same region.

Open pits, underground workings, quarries, waste-rock dumps, tailings facilities and access roads can remove vegetation, alter landforms and change drainage. Associated roads, power infrastructure and settlements may extend impacts beyond the mine boundary. Habitat fragmentation and disturbance are especially concerning in areas of high conservation value or where communities depend directly on local land and water.

B. *Water contamination and tailings*

Risk management must consider both routine releases and low-probability, high-consequence failures. Tailings design should account for site geology, seismic conditions, extreme precipitation, water balance, inspection access and credible failure scenarios. Monitoring should establish baseline conditions before operation and continue after closure where contaminants or altered hydrology may persist.

Mine-related water risks include sediment runoff, altered groundwater flow, metal leaching and acid mine drainage. Where sulfide-bearing minerals react with oxygen and water, acidic drainage may mobilize metals and harm aquatic organisms. Tailings facilities require site-specific design, inspection, emergency planning and long-term maintenance. Some artisanal gold-mining practices use mercury, creating risks of environmental contamination and exposure. The level of risk depends on geology, mineral type, processing, climate and management quality.

C. *Responsible mining and rehabilitation*

Responsible mining begins with baseline studies and avoidance of areas where damage cannot be adequately prevented or remedied. Operators should monitor surface water, groundwater, sediment and air quality; manage tailings and waste rock; reduce process-water demand; and disclose results in accessible formats. Closure plans should be developed and funded from the start, with landform stabilization, soil replacement, revegetation and long-term monitoring. Traceable mineral supply chains and recycling can reduce avoidable demand for virgin extraction, although recycling cannot eliminate the need for all primary minerals.




Fig. 4. Mining life-cycle stages and core environmental risk-management actions.

VIII. ENVIRONMENTAL EFFECTS OF TRANSPORTATION

A. *Climate emissions and energy demand*

Transport emissions depend on system design as much as individual vehicle efficiency. A more efficient vehicle can be offset by longer trips, increased vehicle size or higher total travel demand. Freight policies should assess load factors, empty running, warehousing location and delivery patterns; passenger transport should assess accessibility and service quality, not only average speed.

Road vehicles, aviation, shipping and rail use energy to move people and goods. Fossil-fuel combustion releases carbon dioxide, while additional impacts arise from fuel production, vehicle manufacture, infrastructure construction and maintenance. The footprint varies with mode, distance, occupancy, freight loading, vehicle efficiency and energy source. Electrification can reduce tailpipe emissions, but its full climate benefit depends on electricity generation, battery supply chains and vehicle life-cycle impacts.

B. *Air quality, noise and health*

The distribution of exposure matters: people who live, study or work near busy roads may experience higher pollution and noise than citywide averages suggest. Transport assessments should therefore combine emissions inventories with roadside monitoring, population density, sensitive receptors and travel access. Electric drivetrains reduce exhaust at the point of use but do not eliminate tyre, brake and road-dust particles.

Transport pollution includes exhaust gases and particles as well as non-exhaust emissions from tyres, brakes and road dust. Exposure can be elevated near busy roads, freight corridors, ports and transport hubs. WHO identifies transport-related air pollution as a public-health concern [7]. Noise and vibration can disturb sleep and well-being, while road safety and unequal exposure shape the social consequences of transport systems.

C. *Habitat fragmentation and runoff*

Roads, railways, airports and ports occupy land and may divide habitats into smaller patches. Roads can block wildlife movement, increase collision mortality and introduce noise and light. Runoff can carry oil, metals, salts and particles into waterways. Transport routes can also increase access to

ISSN: 3139-1141

previously remote areas, potentially accelerating settlement, logging, mining or other land conversion.

D. *Sustainable transport responses*

- Build reliable public transport networks and safe walking and cycling infrastructure.
- Support mixed-use planning that shortens everyday trips and improves access to essential services.
- Improve vehicle efficiency, freight logistics, maintenance and loading factors.
- Electrify suitable vehicle fleets alongside cleaner electricity and responsible battery recycling.
- Shift suitable freight to rail and water and improve intermodal connections.
- Use wildlife crossings, ecological corridors and effective drainage where transport routes intersect sensitive habitats.




Fig. 5. Avoid–shift–improve framework for sustainable transport planning.

IX. COMPARATIVE AND CROSS-SECTOR ANALYSIS

The sectors share environmental pathways but differ in their dominant pressures. Agriculture is strongly linked to land, soil, nutrient cycling and water withdrawals. Housing combines land conversion with embodied material impacts and long-term energy use. Industry concentrates process emissions, energy demand and complex waste streams. Mining can create localized but persistent landscape and water risks while supplying materials to other sectors. Transportation contributes to climate forcing and local pollution while shaping land use and ecological connectivity.




Fig. 6. Conceptual relationships between five human activities and shared environmental systems.

A. *Interactions and trade-offs*

A robust comparison should use life-cycle assessment where product footprints are compared, geographic information systems where land and habitat change are assessed, and health-risk assessment where human exposure is the primary concern. These tools answer different questions. Their results become more useful when assumptions, uncertainty and excluded impacts are reported transparently.

Environmental measures can shift rather than eliminate impacts. Electric vehicles may reduce local exhaust emissions but require minerals, manufacturing energy and electricity. Renewable-energy infrastructure can reduce operational emissions while increasing demand for selected metals and land. Dense housing can reduce travel demand but still requires green space, ventilation and heat-resilient design. Agricultural intensification may spare land in some contexts, but it can also increase chemical inputs and harm biodiversity. Life-cycle assessment, cumulative-impact analysis and site-specific ecological thresholds are therefore needed to identify real net benefits.

B. *Environmental justice*

Environmental burdens are distributed unevenly. Agricultural workers, mine workers, low-income communities, informal settlements and residents near industrial facilities or major roads may experience high exposures while having limited influence over decisions. A fair transition requires access to information, meaningful participation, worker protections, affordable clean energy and mobility, and effective grievance mechanisms. Policies should track not only total emissions and resource use but also who receives benefits and who bears risks.

C. *Monitoring indicators*

A practical indicator set should include leading indicators —such as compliance with building codes, nutrient plans, maintenance completion and tailings inspections—as well as outcome indicators such as water quality, habitat condition and measured emissions. Data should be disaggregated by location and affected population where possible, because a favorable average can conceal a severe local hotspot.

Monitoring should combine pressure indicators, environmental-condition measures and social outcomes. Examples include greenhouse-gas emissions per unit of output, land converted or restored, soil organic carbon, freshwater withdrawals, water-quality parameters, particulate matter, habitat connectivity, waste recovery, material intensity, occupational injuries and community exposure. Every indicator should define its unit, baseline year, boundary, data source, uncertainty and reporting frequency. Composite scores should not conceal severe local harms.

X. INTEGRATED RECOMMENDATIONS

A. *Cumulative and life-cycle assessment*

Environmental decisions should follow a mitigation hierarchy: avoid impacts first, minimize those that cannot be avoided, restore affected systems and use offsets only where appropriate and credible. Life-cycle assessment is useful for comparing product and technology footprints; strategic environmental assessment helps evaluate plans and alternatives; and cumulative-impact assessment addresses the

ISSN: 3139-1141

combined pressure from multiple projects. These tools should be complemented by community consultation, baseline data and transparent uncertainty reporting.

B. *Government and regulators*

- Coordinate land-use, climate, water, biodiversity, housing, industry and transport planning.
- Apply environmental and cumulative-impact assessment to major projects before approval.
- Enforce clear standards for air emissions, effluent, waste, building performance and mine closure.
- Use credible monitoring, transparent reporting and proportionate enforcement.
- Protect vulnerable ecosystems and enable meaningful community consultation.

C. *Businesses and project developers*

- Measure direct and supply-chain impacts and disclose methods, boundaries and uncertainty.
- Set time-bound targets backed by investment plans and independently verifiable data.
- Adopt cleaner production, preventive maintenance, water reuse and waste minimization.
- Assess biodiversity, climate risk, worker safety and community exposure throughout project life cycles.
- Avoid unsupported environmental claims and distinguish measured reductions from future intentions.

D. *Communities and individuals*

- Reduce avoidable food waste, conserve water and support soil- and habitat-friendly practices.
- Choose durable and repairable products and reuse materials when safe and practical.
- Improve household energy efficiency and use lower-emissions travel options when accessible.
- Participate in local planning and support transparent, evidence-based environmental decisions.




Fig. 7. Integrated framework for preventing, reducing, controlling and restoring environmental impacts.

XI. IMPLEMENTATION AND EVALUATION FRAMEWORK

A practical implementation cycle begins with a baseline inventory, identifies high-impact activities and exposed populations, selects prevention and mitigation measures, assigns responsibility and financing, and then evaluates performance against a defined timeline. The cycle should be repeated as new monitoring data become available. Baselines should cover environmental conditions before a project begins, while indicators should be selected according to the decision being made rather than availability alone.

XII. RESEARCH GAPS AND FUTURE DIRECTIONS

Research needs include comparable datasets for regions and supply chains, especially where informal activities and imported goods are poorly represented. More studies should connect global life-cycle footprints with local ecological thresholds, including watershed limits and biodiversity sensitivity. Monitoring should better represent cumulative exposure among communities affected by multiple pollution sources. Transition technologies should be assessed across their full life cycle so that reductions in one impact category do not hide increases in another. Participatory research is also needed to understand how environmental costs, employment changes and benefits are distributed.

Future studies could apply a systematic-review protocol with documented databases, search strings, inclusion criteria and screening counts. Quantitative comparisons could use defined functional units such as one tonne of crop output, one square metre-year of housing, one tonne of manufactured material, one tonne of mineral produced or one passenger-kilometre of travel. These comparisons require consistent system boundaries and explicit uncertainty analysis. Local case studies could complement global assessments by measuring water quality, air pollution, land-cover change and community experience.

XIII. LIMITATIONS

This paper synthesizes selected authoritative literature and is not an exhaustive systematic review of all global publications. Its broad sectoral descriptions cannot represent every agricultural system, building type, industrial process, mineral deposit or transport network. Estimates differ by reporting year, method and system boundary, so figures from different sources should not be added or compared as though they were calculated identically. The paper reports no new experimental data or quantitative effect size. Before submission, each reference and citation should be verified against the original publication and the final manuscript should be checked using the journal's required similarity-detection process.

XIV. CONCLUSION

Agriculture, housing, industry, mining and transportation are essential to modern societies, yet poorly planned or resource-intensive practices can contribute to climate change, habitat loss, soil degradation, freshwater stress, air and water pollution, waste generation and risks to human health. Their effects are interconnected: housing relies on industrial materials and mined minerals; transport infrastructure reshapes land use; agricultural supply chains depend on energy and logistics; and industry and mining can affect the same watersheds and communities.

The literature supports a shift from isolated pollution control toward integrated life-cycle management. Priority measures include protecting ecosystems, improving

ISSN: 3139-1141

agricultural soil and water practices, building efficient and resilient housing, cleaning industrial processes, strengthening mining safeguards and rehabilitation, and providing safe, accessible low-emissions transport. Effective governance, transparent monitoring, community participation and fair distribution of transition costs are essential. Sustainable development does not require abandoning food production, housing, manufacturing, mineral supply or mobility; it requires delivering these services with less resource use, fewer emissions, reduced ecological harm and stronger accountability.

ACKNOWLEDGMENT

The authors acknowledge the publicly available scientific assessments and institutional resources cited in this review. No external funding or specific institutional grant is claimed in this manuscript.

REFERENCES

[1] Intergovernmental Panel on Climate Change, “Climate change 2023: Synthesis report,” IPCC, Geneva, Switzerland, 2023. [Online]. Available: https://www.ipcc.ch/report/ar6/syr/

[2] Intergovernmental Panel on Climate Change, “Climate change 2022: Mitigation of climate change,” Working Group III contribution to the Sixth Assessment Report, Cambridge University Press, 2022. [Online]. Available: https://www.ipcc.ch/report/ar6/wg3/

[3] Food and Agriculture Organization of the United Nations, “The state of the world’s land and water resources for food and agriculture 2025: The potential to produce more and better,” FAO, Rome, 2025. [Online]. Available: https://www.fao.org/publications/fao-flagship-publications/the-state-of-the-worlds-land-and-water-resources-for-food-and-agriculture/

[4] United Nations Environment Programme and Global Alliance for Buildings and Construction, “Global status report for buildings and construction 2024/2025: Not just another brick in the wall,” UNEP, 2025. [Online]. Available: https://wedocs.unep.org/handle/20.500.11822/47214

[5] United Nations Environment Programme, “Artisanal gold mining and biodiversity (ASGM): Restoring the balance,” UNEP, 2023. [Online]. Available: https://wedocs.unep.org/handle/20.500.11822/43323

[6] United Nations Environment Programme, “Chemical pollution and biodiversity,” UNEP. [Online]. Available: https://www.unep.org/en/chemical-pollution-and-biodiversity

[7] World Health Organization, “Transport: Sectoral solutions for air pollution and health,” WHO, July 17, 2025. [Online]. Available: https://www.who.int/publications/i/item/B09368

[8] United Nations Environment Programme and Global Alliance for Buildings and Construction, “Global status report for buildings and construction 2025–2026: Building fast. Falling short,” UNEP, 2026. [Online]. Available: https://wedocs.unep.org/handle/20.500.11822/49531